



# SELINEXOR

Antineoplastic, XPO1 inhibitor

Availability <sup>1</sup>	Brand name	Quick guide
Tablet, film-coated 20 mg	Xpovia	 
<b>Effect of food</b>	Food does not significantly affect absorption. <sup>1</sup> Giving the dose within 30 minutes of food or enteral nutrition may help with nausea. <sup>2</sup>	
<b>Warnings</b>	<b>Hazard.<sup>1</sup> Do not crush the tablet.</b> Occupational exposure may be harmful. <b>Do not crush or disperse the tablet if you are pregnant.</b> Contact a pharmacist or a medicines information service for advice.	
<b>Other</b>	May cause dry mouth. <sup>1</sup>	

## What to do for people with enteral feeding tubes

### **Do not crush or break the tablet.<sup>1</sup>**

Hospital pharmacies can prepare a 5 mg/mL suspension in ORA-Blend that is suitable for nasogastric and gastrostomy tubes.<sup>2</sup>

Wear gloves when measuring and giving the dose. Contact with the skin causes irritation and a rash.<sup>2</sup>

Gently invert the bottle to re-suspend the contents. Do not shake.<sup>2</sup>

## What to do for people with swallowing difficulties

### **Do not crush or break the tablet.<sup>1</sup>**

Hospital pharmacies can prepare a 5 mg/mL oral suspension in ORA-Blend.<sup>2</sup>

Wear gloves when measuring and giving the dose. Contact with the skin causes irritation and a rash.<sup>2</sup>

Gently invert the bottle to re-suspend the contents. Do not shake.<sup>2</sup>

## REFERENCES:

1. Product information. Available from [www.tga.gov.au](http://www.tga.gov.au). Accessed 07/10/2022.

2. Medical information. Selinexor preparation as a suspension for oral, nasogastric tube or gastrostomy tube administration (2022JUL22v1.0). Melbourne: Antengene Aus; 22/07/2022.