ENCORAFENIB

Antineoplastic, BRAF inhibitor

Availability ¹		Brand name	Quick guide
Capsule 50 mg,	75 mg	Braftovi	
Effect of food	Food does not significantly affect absorption. Can be given without regard to food. ¹ Avoid grapefruit and grapefruit juice. ¹		
Warnings	Hazard . ¹ Occupational exposure may be harmful. Contact a pharmacist or a medicines information service for advice.		
Other	May cause impaired or altered sense of taste. ¹		
What to do for people with enteral feeding tubes			
	See <i>Warnings</i> . There is no information about the safety or efficacy of administration by enteral feeding tube. Contact a pharmacist or a medicines information service for advice, or contact the prescriber to consider other options.		

What to do for people with swallowing difficulties

Do not crush or break the tablet.

See Warnings.

Open the capsule(s) and mix the contents with a tablespoon of apple puree. Stir for 2 minutes to ensure a smooth consistency.²

Up to 6 capsules can be mixed with 20 mL of apple puree.²

Stable for 2 hours at room temperature.²

REFERENCES:

Product information. Available from www.tga.gov.au, Accessed 20/01/2023.
Medical and patient consumer department. Information Request - Binimetinib (Mektovi) - SHPA Don't Rush to Crush (DRTC) Monograph Review [email]. North Sydney: Pierre Fabre Group; 14/12/2022.