## **SEMAGLUTIDE**

BRAND NAME OZEMPIC

DRUG CLASS Glucagon-like peptide-1 (GLP-1) agonist

AVAILABILITY Prefilled pen contains 2 mg/1.5 mL (0.25 mg or 0.5 mg dose) or 4 mg/3 mL (1 mg dose)

of semaglutide. Also contains dibasic sodium phosphate dihydrate, propylene glycol,

phenol, hydrochloric acid and sodium hydroxide.1

The solution is clear and colourless.1

pH 7.4<sup>1</sup>

PREPARATION Ready to use

STABILITY Store at 2 to 8 °C. Do not freeze. Protect from light.1

Once in use the pen is stable for 6 weeks below 30 °C or at 2 to 8 °C. 1

Remove and discard the needle after each dose and store the pen with the cap on to

protect from light.1

**ADMINISTRATON** 

**IM** injection Contraindicated<sup>1</sup>

**SUBCUT injection** Inject into the abdomen, thigh or upper arm. Suitable for self-administration after

appropriate patient education.1

Can be used with needles that are up to 8 mm in length. The pen is designed to be

used with NovoFine disposable needles.1

IV injection Contraindicated<sup>1</sup>
IV infusion Contraindicated<sup>1</sup>

COMPATIBILITY Do not mix with fluids or other medicines.<sup>1</sup>

INCOMPATIBILITY No information

SPECIAL NOTES The dose is given once weekly on the same day each week.<sup>1</sup>

Anaphylactic reactions have been reported.1

## **REFERENCES**

<sup>1.</sup> Product information. Available from www.tga.gov.au. Accessed 14/07/2020.