LEFLUNOMIDE

Disease modifying anti-rheumatic drug (DMARD)

Quick guide Availability1 Brand name Tablet, film-coated 10 mg, 20 mg Arava, Ataris, Lunava Generics are available

Effect of food Can be given without regard to food.1

Hazard.^{1,2} Occupational exposure may be harmful. Wear a mask and gloves if dispersing Warnings

the tablet. Teratogen.^{1,2} Do not disperse the tablet if you are pregnant. Contact a

pharmacist or a medicines information service for advice.

Other May cause dry mouth, mouth ulcers and an impaired or altered sense of taste.¹

What to do for people with enteral feeding tubes

Option 1 Do not break or crush the tablet.

See Warnings. Wear a mask and gloves.

Disperse the tablet in 10–20 mL of water in an enteral syringe. Shake gently, the tablet will disperse within 2 to 7 minutes.³

See Method A for instructions.

What to do for people with swallowing difficulties

Option 1 Do not break or crush the tablet.

See Warnings. Wear a mask and gloves. Use an oral dispenser.

Disperse the tablet in water.

Shake gently, the tablet will disperse in 10–20 mL of water in 2 to 7 minutes.³

See Method D for instructions.

If the person cannot swallow thin fluids, contact a pharmacist or a medicines information Option 2

service or the prescriber for advice.

REFERENCES:

Product information. Available from www.tga.gov.au. Accessed 06/10/2022.
National Institute for Occupational Safety and Health. NIOSH list of antineoplastic and other hazardous drugs in healthcare settings 2020. Cincinnati, OH: Department of Health and Human Services; 2020.
SHPA. Drug dispersion database; 2015, 2017.